

(19)



Europäisches Patentamt
European Patent Office
Office européen des brevets



(11)

EP 0 659 344 B1

(12)

EUROPEAN PATENT SPECIFICATION

(45) Date of publication and mention
of the grant of the patent:
04.07.2001 Bulletin 2001/27

(51) Int Cl.7: **A21D 8/04**, A21D 2/22,
C12G 3/02

(21) Application number: 94203629.4

(22) Date of filing: 14.12.1994

(54) **Dry yeast compositions**

Trockenhefe Zusammensetzungen

Compositions de levure séché

(84) Designated Contracting States:
**AT BE CH DE DK ES FR GB GR IE IT LI LU MC NL
PT SE**

(30) Priority: 24.12.1993 EP 93203697

(43) Date of publication of application:
28.06.1995 Bulletin 1995/26

(60) Divisional application:
01200011.3 / 1 090 553

(73) Proprietor: **DSM N.V.**
6411 TE Heerlen (NL)

(72) Inventor: **Groenendaal, Jan Willem**
NL-2611 EE Delft (NL)

(74) Representative:
Matulewicz, Emil Rudolf Antonius, Dr. et al
DSM Patents & Trademarks
Office Delft
P.O. Box 1
2600 MA Delft (NL)

(56) References cited:
EP-A- 0 310 460 EP-A- 0 616 030
EP-A- 0 619 947 FR-A- 1 509 676
FR-A- 2 233 941 FR-A- 2 690 813
US-A- 3 843 800

- Römpp Chemie Lexikon, 9th ed., Georg Thieme Verlag, Stuttgart, 1990, volume 2, p. 1641
- List, P. H., Arzneiformlehre, MVG, Stuttgart, 1985, p. 82 - 88

Note: Within nine months from the publication of the mention of the grant of the European patent, any person may give notice to the European Patent Office of opposition to the European patent granted. Notice of opposition shall be filed in a written reasoned statement. It shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

EP 0 659 344 B1

Description

[0001] The present invention relates to dry yeast compositions, the production thereof and their use in bakery products and beverages.

[0002] The manufacture of a yeast composition starts with a small sample of a pure culture. This sample is used to inoculate the first of a series of fermentations in fermenters of successively increasing size. The first few are mildly aerated batch fermentations. Only the last two (or sometimes three) stages are performed using full aeration and incremental feeding of molasses. These fedbatch fermentations are carried out in fermenters having a volume of 100m³ or more. Fermentation is typically carried out for a total of 12-20 hours during which time about 10,000-30,000 kg of compressed yeast is produced.

[0003] Further processing includes separating the yeast from the broth by centrifugation and washing which results in a yeast cream (17-23% (w/w) dry matter content).

[0004] The yeast cream may be processed into compressed yeast (27-33% (w/w) dry matter content) which is either sold as such or extruded and dried to produce active dry yeast (ADY) or instant dry yeast (IDY) with moisture contents of 6-8% (w/w) and 2-8% (w/w), respectively.

[0005] In the case of ADY, drying usually takes place in belt or rotolouvre (drum) dryers. For IDY production, fluidized-bed drying is commonly used. Drying of the yeast to a level of about 20% w/w water content involves only the evaporation of free water. Further reduction of the moisture content requires removal of a portion of the bound water from the yeast which may cause damage to the yeast cell membrane. US patents 3,843,800 and 4,248,420 describe use in such a drying process of a wetting agent such as a glycerol ester and/or fatty acid ester of propylene glycol so as to preserve the desired high direct leavening activity of the yeast.

[0006] Dry yeast loses part of its leavening activity during both drying and rehydration. Dry yeasts are still commonly used in the bakery trade because of their extended stability and because refrigeration is unnecessary. Dry yeasts are used in wine making to obtain a fast and reproducible fermentation thereby avoiding the risk of failure of natural fermentation. Moreover, the yeast is immediately available throughout the year.

[0007] Instant dry yeast (IDY) is the latest type of baker's yeast, which was introduced in the early 1970s (see for example US patent 3,843,800). This was followed a few years later by introduction of instant dry wine yeast (IWY), which can be regarded as a special form of instant dry yeast. To obtain a high quality IDY, compressed yeast of relatively high protein content (42-60% (w/w)) must be dried in a quick drying process. The leavening activity of IDY under conditions of application is about 75-85% that of compressed yeast. The shelf life in a vacuum-sealed package is comparable to that of ADY.

[0008] IDY is presented typically in the form of very small rods that are highly porous and easy to rehydrate. On the one hand, this allows immediate use, without prior rehydration. On the other hand, the high porosity gives easy access to water and oxygen (from air) which results in a rather rapid loss of activity upon exposure to atmospheric conditions. For satisfactory results, IDY should be used within 3-5 days of opening the package. Moreover, the high porosity of IDY makes it sensitive to extreme rehydration conditions.

[0009] IDY usually has a moisture content of 2-8% (w/w) and a protein content between 42 and 60% (w/w) on a dry matter basis.

[0010] In baking, besides bakers' yeast, processing aids such as bread improvers are used, including oxidising and reducing agents, enzymes such as redox enzymes, α -amylases, amyloglucosidases, hemicellulases, cellulases and proteases, lipases and phospholipases, emulsifiers and fatty materials. Yeast, enzymes and redox agents are added separately to doughs. Yeast may be added in cream, compressed, active dry or instant dry form. Enzymes may be added in dry powder or in dissolved form. Redox agents are in most cases used in powder form.

[0011] Separate weighing and dosing of these various ingredients increases the number of actions which have to be performed by the operators of the production process. Inherent in this increased number of actions is a greater chance of introducing errors resulting in negative impact on the quality of the end product. Moreover, working with dusty materials may initiate allergic reactions.

[0012] Mixing of ingredients with dry granular yeast or instant dry yeast may result in homogeneous products directly after mixing. However, during transport and storage prior to use, this type of product tends to lose homogeneity (see Example 1). To solve these problems various solutions have been proposed.

[0013] J-73040748 describes the mixing of granular semi-dry yeast (moisture content of 35-45% w/w) with a wheat flour improving agent for use in breadmaking. In such a mixture, the stability of both the flour improving agent and yeast is very limited due to the high water content. Thus, special attention has to be paid to storage and transport conditions.

[0014] DE-2515029 describes the production of active dry yeast by vacuum drying and coformulation of the dry yeast with spray dried malt extract or maltodextrins. Malt extract or maltodextrins are added as a dewatering agent. However, this vacuum drying technique cannot be applied economically on a commercial scale due to unacceptable loss in leavening activity. In general, a yeast composition produced by this technique will be in powder or dust form, which

may result in allergy.

[0015] The present invention provides a dry yeast composition which has the advantage of provision of dry yeast with a bread-improving agent, but which has none of the above-mentioned disadvantages such as dust formation and loss of homogeneity of the composition during transport.

5 [0016] Thus, in one aspect the present invention provides a dry granular yeast composition having a moisture content of not more than about 8% (w/w) and containing about 0.1 to 8% (w/w) of a bread-improving agent, wherein the dry yeast is present in granular form and said bread-improving agent is present as a coating on the dry yeast granules in the form of adhered particles.

[0017] In principle, a dry yeast composition of the present invention may have substantially the same granule size or not a greatly increased granule size compared to the starting dry yeast before the bread-improving agent is added. Additionally, a composition of the invention has the same convenience of use as dry yeast, which makes application possible without adapting operational procedures or equipment.

[0018] Preferably, the bread-improving agent is added to the dry yeast at 1 to 4% (w/w). The bread-improving agent may preferably comprise one or more enzymes and/or ascorbic acid. Enzymes for this purpose can be selected from carbohydrases such as α -amylase, amyloglucosidase, hemicellulase, cellulase and glucanase, protein modifying enzymes such as proteases and peptidases, redox enzymes such as glucose oxidase, sulfhydryloxidase and lipoxidase (lipoxigenase), peptidyl transferases such as γ -glutamyl transferase and lipid modifying enzymes such as lipases and phospholipases.

[0019] The bread-improving agent is coated on to the dry yeast granules. The coating may be in the form of very small adhered particles. If a bread-improving agent is employed to provide a particle coating, preferably at least 50% of the particles will have a size smaller than 50 μ m. It has been found advantageous for at least 80% of the particles to have a smaller size than 50 μ m. Particles of a bread-improving agent which are too large for direct use in a composition of the invention can be reduced to appropriate size using suitable equipment known in the art such as a milling apparatus.

[0020] The bread-improving agent particles are stuck on to the dry yeast using suitable adhesives. These will be in general foodgrade adhesives, preferably have a keeping quality of at least two years when applied on dry yeast and will not influence the taste or flavour of the dry yeast, the dough prepared with the yeast or the final bread product. The adhesive is preferably added to a mixture of yeast granules and microfine bread-improving particles, e.g. at 0.5 to 1.0% (w/w). During application of the adhesive, continuous mixing is preferably carried out. The adhesive is preferably slowly supplied either continuously or batch-wise using small portions each time. After completion of addition of the adhesive, the mixing will continue until substantially all the bread-improving particles are stuck on to the dry yeast granules. While the adhesive can be poured on to the mixture, preferably the adhesive is sprayed on to the mixture to obtain a more uniform distribution. An in-line mixing process may be used (see Example 6).

[0021] Suitable adhesives are e.g. oils such as soy oil, cotton seed oil, rape seed oil, sunflower oil, corn oil, peanut oil, olive oil, paraffin oil, triglycerides, liquid fats and mixtures thereof. Fractionated oils can be used. The adhesive may include one or more additives which are beneficial in improving the sticking characteristics. Thus, for example, lecithin may be advantageously mixed with soy oil.

[0022] In further aspects, the present invention additionally provides use of a dry granular yeast composition of the invention for incorporation into a dough or for fermentation of a beverage and doughs and beverage compositions incorporating such a dry yeast composition.

[0023] The following examples illustrate the invention.

Example 1 (comparative example)

45 [0024] 2,700 g Femipan™ (dry yeast of Gist-brocades) were homogeneously mixed with 36 g ascorbic acid, 6 g fungal α -amylase Femizyme™ P₂₀₀ (Gist-brocades, 4740 PU/g) and 48g hemicellulase Femizyme™ H₁₀₀₀ (Gist-brocades, hemicellulase activity 13,500 HU/g and α -amylase activity 942 PU/g) in a Hobart mixer. Directly after mixing, portions of 450 g were weighed and packed in aluminium bags, which were closed at reduced pressure.

[0025] The homogeneity of the contents of three packs was tested directly after packing by opening each pack at three places, near the top, in the middle, and near the bottom, and withdrawing samples of 25 g from each opening. In these samples, the levels of ascorbic acid, fungal α -amylase and hemicellulase were analyzed according to the following methods:

- ascorbic acid analysis was carried out according to the conventional method of Boehringer.
- 55 - fungal α -amylase activity was determined using Phadebas™ tablets from Pharmacia. In this method, solubilization of dye-labelled starch by α -amylase over 15 minutes in a buffer at pH 5.5 and 30°C is measured spectrophotometrically. α -amylase activity is expressed in Phadebas Units (PU) using an Aspergillus oryzae fungal α -amylase preparation of 10,000 PU/g as an internal standard. One Phadebas Unit defined in this way equals about 10 SKB

units, used in the baking industry.

- fungal hemicellulase activity was determined by measuring the amount of reducing sugars produced over a pre-determined time period in the micro-assay as described by Leathers T.D., Kurtzman, C.P. and Detroy, R.W. in Biotechnol. Bioeng. Symp., (1984) 14, 225. In this paper, the hemicellulase unit (HU) is also defined.

[0026] The results are summarised in Table 1.

[0027] Another three packs prepared as described above were stored for two weeks in a refrigerator at 4°C. Afterwards, these packs were placed in a conventional carton for instant yeast packs and surrounded by conventional instant yeast packs. This case was transported by a heavy goods vehicle over about 2500 km such that the packs were in a vertical position. Afterwards, the three test packs were stored in a refrigerator again for another four weeks. The homogeneity was then tested in the above described way. The analysis results are summarised in Table 1.

TABLE 1

| Homogeneous mix | | | Relative Amount recovered (%) | |
|-----------------|-------------------|-----------|-------------------------------|-----------------|
| | | | directly | after transport |
| Top | ascorbic acid | 0.013 g/g | 101.5 | 85.2 |
| | α -amylase | 26 PU/g | 98.5 | 91.4 |
| | hemicellulase | 232 HU/g | 100.8 | 92.5 |
| Middle | ascorbic acid | 0.013 g/g | 102.0 | 89.2 |
| | α -amylase | 26 PU/g | 98.1 | 94.6 |
| | hemicellulase | 232 HU/g | 99.6 | 91.4 |
| Bottom | ascorbic acid | 0.013 g/g | 99.1 | 112.3 |
| | α -amylase | 26 PU/g | 101.7 | 108.6 |
| | hemicellulase | 232 HU/g | 98.7 | 106.9 |

[0028] From these results, it is clear that during storage and transport, the mixture of yeast, ascorbic acid and enzymes lost homogeneity.

Example 2

Mixing process for 1 kg dry yeast composition

[0029] a) The following components were weighed :

| | | |
|---|----------------------|-----------------|
| - Fermipan™ (dry yeast of Gist-brocades) | 975.700 g = | 97.57 % |
| - Vitamin C microfine | 16.873 g = | 1.69 % |
| - Hemicellulase 25,690 HU/g | 0.856 g = | 0.08 % |
| - α -amylase 11,400 PU/g | 1.632 g = | 0.16 % |
| - Kriskol 3000 (fractionated fat, Loders, Crokiaan) | 5.000 g = 1,000 g | 0.50 % 100 % |

b) A premix was prepared consisting of 100 g Fermipan™ combined with the total amount of Vitamin C and enzymes by mixing the components with a spoon in a 250ml beaker.

c) The premix was put in the mixing beaker of a Hobart planetary mixer together with the remaining part of the Fermipan.

d) Mixing was started and the Kriskol 3000 was added in 30 sec. The total mixing time was 10 mins.

e) The final product was vacuum packed in aluminium bags and stored at 5°C.

Example 3**Mixing process for 30 kg dry yeast composition**

5 [0030]

a) The following components were weighed:

10

| | | |
|---|---------------------|-----------------|
| - Fermipan™ | 29,121 g = | 97.07 % |
| - Vitamin C microfine | 506.2 g = | 1.69 % |
| - Hemicellulase 25,690 HU/g | 25.7 g = | 0.08 % |
| - α -amylase 11,400 PU/g | 49.0 g = | 0.16 % |
| - Durkex 500 (mixture of soy oil and cotton seed oil, Otto Aldag, Hamburg) | 300 g = 30,000 g | 1.00 % 100 % |

15

b) A premix was prepared by combining 4,367g Fermipan™, the total amount of Vitamin C and enzymes and 99g of Durkex 500 in a Hobart planetary mixer.

20

c) The premix was put in a Nauta conical mixer together with the remaining part of the Fermipan.

d) Mixing was started and the remaining part of the Durkex 500 was added in 40 sec. The total mixing time was 10 mins.

e) The final product was vacuum packed in aluminium bags and stored at 5°C.

25

Example 4 (not part of the claimed invention)**Granulating process for 500 kg of Vitamin C and enzymes prepared in a Multi-Stage spray-dryer and mixing process to produce 10,000 kg of dry yeast composition.**

30

[0031]

a) The following components were weighed:

35

| | | |
|---------------------------------|------------|--------|
| - Vitamin C | 15.6 kg = | 3.12% |
| - Vitamin C sodium | 420.2 kg = | 84.04% |
| - Hemicellulase 25,690 HU/g | 22.1 kg = | 4.42% |
| - α -amylase 11,400 PU/g | 42.1 kg = | 8.42% |
| | 500 kg = | 100 % |
| - water 750 kg | | |

40

b) A solution was prepared by mixing the Vitamin C and enzymes with the water in a vessel of 1500 l equipped with a turbine stirrer.

45

c) Directly after preparation of the solution, it was fed to a Stork Multi-Stage spray-dryer and dried with fine return at an inlet temperature of about 160°C and an outlet temperature of about 90°C.

d) 197 kg of the MSD-granulate was transported to a conical Nauta mixer with a capacity of 15 m³ together with 9,803 kg of Fermipan™. The total mixing time was 20 mins.

e) The dried product was vacuum packed in aluminium bags and stored at 5°C.

50

Example 5 (not part of the claimed invention)**Coating process for 5 kg dry yeast composition**

55

[0032]

a) The following components were weighed:

| | | |
|---------------------------------|--------------------|-----------------|
| - Fermipan™ | 4,803.5 g = | 96.7 % |
| - Vitamin C microfine | 3 g = | 0.06 % |
| - Vitamin C sodium | 81.4 g = | 1.63 % |
| - Hemicellulase 25,690 HU/g | 4.3 g = | 0.08 % |
| - α -amylase 11,400 PU/g | 8.2 g = | 0.16 % |
| - hydroxypropyl cellulose | 100 g = 5,000 g | 2.00 % 100 % |
| - water 1,979 g | | |

b) A solution was prepared by mixing the Vitamin C, enzymes and hydroxypropyl cellulose with the water in a 3 l vessel equipped with a turbine stirrer.

c) Directly after the preparation of the solution, it was fed at a rate of about 15 g per min to an Aeromatic MG-1 fluidized bed coating apparatus equipped with a Wurster column containing the Fermipan™. The inlet temperature was about 75°C and the outlet temperature about 45°C.

d) The dried product was vacuum packed in aluminium bags and stored at 5°C.

Example 6

In-line mixing process for 10,000 kg dry yeast composition

[0033]

a) The following components were weighed:

| | | |
|---------------------------------|-----------------------|-----------------|
| - Fermipan™ | 9,706.4 kg = | 97.07 % |
| - Vitamin C microfine | 168.7 kg = | 1.69 % |
| - Hemicellulase 25,690 HU/g | 8.6 kg = | 0.08 % |
| - α -amylase 11,400 PU/g | 16.3 kg = | 0.16 % |
| - Durkex 500 | 100 kg = 10,000 kg | 1.00 % 100 % |

b) A premix was prepared of 1,456 kg of Fermipan™ combined with the total amount of Vitamin C and enzymes and 33 kg of the Durkex 500 in a Nauta conical mixer. The mixing time was 20 mins.

c) This premix was fed with the aid of pneumatic transport to an in-line mixer together with the remaining part of the Fermipan™ and the remaining part of the Durkex 500. The flow of Fermipan™, the premix and the Durkex 500 were adjusted to accord with the above-mentioned composition with the aid of automatic dosing units.

d) Directly following the in-line mixer, the product was vacuum packed in aluminium bags and stored at 5°C.

Claims

1. A dry granular yeast composition having a moisture content of not more than about 8% (w/w) and containing about 0.1 to 8% (w/w) of a bread-improving agent, wherein the dry yeast is present in granular form, and wherein the bread-improving agent comprises one or more enzymes and/or ascorbic acid and is present as a coating on the dry yeast granules in the form of particles adhered to said granules by an adhesive.
2. A composition as claimed in claim 1, wherein at least 50% preferably at least 80%, of said adhered particles have a diameter of less than 50 μ m.
3. A composition as claimed in claim 1 or 2, wherein said bread-improving agent is present in the form of particles adhered to the dry yeast by means of an adhesive comprising one or more edible oils, triglycerids and liquid fats.

4. A composition as claimed in any one of the preceding claims, wherein the bread-improving agent comprises at least one of a hemicellulase and an α -amylase.
- 5 5. A process for the production of a composition as claimed in any one of the claims 1 to 4, wherein dry yeast granules having a moisture content of not more than about 8% (w/w) are mixed with smaller particles of a bread-improving agent in the presence of an adhesive so that said yeast granules become coated with particles of the bread-improving agent.
- 10 6. Use of a composition as claimed in any one of the claims 1 to 4 for incorporation into a dough or for fermentation of a beverage.

Patentansprüche

- 15 1. Granulierte Trockenhefezusammensetzung mit einem Feuchtigkeitsgehalt von nicht mehr als etwa 8% (Gew/Gew) und einem Gehalt von etwa 0,1 bis 8% (Gew/Gew) eines Brotverbesserungsmittels, worin die Trockenhefe in Granulatform vorliegt sowie das Brotverbesserungsmittel ein oder mehrere Enzyme und/oder Ascorbinsäure beinhaltet sowie als Beschichtung auf dem Trockenhefegranulat in Form von Teilchen, die mittels eines Klebstoffs an dem Granulat haften, vorhanden ist.
- 20 2. Zusammensetzung nach Anspruch 1, worin mindestens 50%, vorzugsweise mindestens 80%, der anhaftenden Teilchen einen Durchmesser von weniger als 50 μ m aufweisen.
- 25 3. Zusammensetzung nach Anspruch 1 oder 2, worin das Brotverbesserungsmittel in Form von Teilchen vorliegt, die mittels eines Klebstoffs an der Trockenhefe haften, der ein oder mehrere Speiseöle, Triglyceride und Flüssigfette enthält.
- 30 4. Zusammensetzung nach einem der vorstehenden Ansprüche, worin das Brotverbesserungsmittel mindestens einen der Stoffe Hemicellulase und α -Amylase enthält.
- 35 5. Verfahren zum Herstellen einer Zusammensetzung gemäß einem der Ansprüche 1 bis 4, worin Trockenhefegranulat mit einem Feuchtigkeitsgehalt von nicht mehr als etwa 8% (Gew/Gew) mit kleineren Teilchen eines Brotverbesserungsmittels in Gegenwart eines Klebstoffs derart gemischt werden, daß das Hefegranulat mit den Teilchen des Brotverbesserungsmittels beschichtet wird.
6. Verwendung einer Zusammensetzung nach einem der Ansprüche 1 bis 4 zur Einarbeitung in einen Teig oder zur Fermentation eines Getränks.

Revendications

- 45 1. Composition de levure granulaire sèche ayant une teneur en humidité qui n'est pas supérieure à environ 8% (p/p) et contenant environ 0,1 à 8% (p/p) d'un agent d'amélioration du pain, dans laquelle la levure sèche est présente sous forme granulaire, et dans laquelle l'agent d'amélioration du pain comprend un ou plusieurs enzymes et/ou de l'acide ascorbique et est présent sous la forme d'un enrobage sur les granulés de levure sèche sous la forme de particules collées auxdits granulés par un adhésif.
- 50 2. Composition selon la revendication 1, dans laquelle au moins 50%, de préférence au moins 80%, desdites particules collées ont un diamètre inférieur à 50 μ m.
3. Composition selon la revendication 1 ou 2, dans laquelle ledit agent d'amélioration du pain est présent sous la forme de particules collées à la levure sèche au moyen d'un adhésif comprenant une ou plusieurs huiles comestibles, des triglycérides et des graisses liquides.
- 55 4. Composition selon l'une quelconque des revendications précédentes, dans laquelle l'agent d'amélioration du pain comprend au moins une hémicellulase et une α -amylase.
5. Procédé de production d'une composition selon l'une quelconque des revendications 1 à 4, dans lequel des gra-

EP 0 659 344 B1

nulés de levure sèche ayant une teneur en humidité qui n'est pas supérieure à environ 8% (p/p) sont mélangés avec de plus petites particules d'un agent d'amélioration du pain en présence d'un adhésif de telle sorte que lesdits granulés de levure soient enrobés de particules de l'agent d'amélioration du pain.

- 5 6. Utilisation d'une composition selon l'une quelconque des revendications 1 à 4 pour incorporation à une pâte ou pour fermentation d'une boisson.

10

15

20

25

30

35

40

45

50

55